
Rules for Means of Random Variables

Rule 1
 If X is a random variable and a and b are fixed numbers, then

$$\mu_{a+bX} = a+b\mu_X$$

Rule 2
 If X and Y are random variables, then

$$\mu_{X+Y} = \mu_X + \mu_Y$$

Statistics 528 - Lecture 17
 Prof. Kate Calder 1

Variance of a Discrete Random Variable

Suppose that X is a discrete random variable whose distribution is

Value of X		x_1	x_2	x_3	...	x_k
Probability		p_1	p_2	p_3	...	p_k

and that μ_X is the mean of X. The **variance** of X is

$$\sigma^2_X = (x_1 - \mu_X)^2 p_1 + (x_2 - \mu_X)^2 p_2 + \dots + (x_k - \mu_X)^2 p_k$$

$$\Sigma (x_i - \mu_X)^2 p_i$$

The **standard deviation** of X is the square root of the variance.

Statistics 528 - Lecture 17
 Prof. Kate Calder 2

Example: Free-Throw (Lecture 16)
 What is the standard deviation of shots made by a 58% shooter who shoots three shots?

The probability distribution for X (number of shots made):

x	P(X=x)
0	0.08
1	0.1+0.1+0.1=0.3
2	0.14+0.14+0.14 = 0.42
3	0.2

Statistics 528 - Lecture 17
 Prof. Kate Calder 3

From lecture 17, $\mu_X = 1.74$

$$\sigma^2_X = (0-1.74)^2 P(X=0) + (1-1.74)^2 P(X=1) + (2-1.74)^2 P(X=2) + (3-1.74)^2 P(X=3)$$

$$= (3.0276)(0.08) + (0.5476)(0.3) + (0.0676)(0.42) + (1.5876)(0.2)$$

$$= 0.7524$$

$$\sigma_X = \text{sqrt}(\sigma^2_X) = 0.8674$$

Statistics 528 - Lecture 17
 Prof. Kate Calder 4

Rules for Variances

Rule 1
 If X is a random variable and a and b are fixed numbers, then

$$\sigma^2_{a+bX} = b^2 \sigma^2_X$$

Rule 2
 If X and Y are *independent* random variables, then

$$\sigma^2_{X+Y} = \sigma^2_X + \sigma^2_Y$$

$$\sigma^2_{X-Y} = \sigma^2_X + \sigma^2_Y$$

This is the **addition rule for variances of independent random variables.**

Statistics 528 - Lecture 17
 Prof. Kate Calder 5

Rule 3
 If X and Y are dependent and have correlation ρ , then

$$\sigma^2_{X+Y} = \sigma^2_X + \sigma^2_Y + 2\rho\sigma_X\sigma_Y$$

$$\sigma^2_{X-Y} = \sigma^2_X + \sigma^2_Y - 2\rho\sigma_X\sigma_Y$$

This is the **general rule for variances of random variables.**


Statistics 528 - Lecture 17
 Prof. Kate Calder 6

General Probability Rules

General Addition Rule for Unions of Two Events
For any two events A and B,

$$P(A \text{ or } B) = P(A) + P(B) - P(A \text{ and } B)$$

Statistics 528 - Lecture 17
Prof. Kate Calder 7



Statistics 528 - Lecture 17
Prof. Kate Calder 8

What is P(A and B)?

- If A and B are *independent*, then by the multiplication rule
 $P(A \text{ and } B) = P(A)P(B)$
- If A and B are *dependent*, then we need a general multiplication rule:
 $P(A \text{ and } B) = P(A)P(B|A)$
where $P(B|A)$ is the *conditional probability* that B occurs given the information that A occurs.

Statistics 528 - Lecture 17
Prof. Kate Calder 9

- Definition of **independence**: Two events A and B are independent if
 $P(A|B) = P(A)$ and $P(B|A) = P(B)$.

So our old multiplication rule for independent events is a special case of the general multiplication rule.

- Definition of **conditional probability**: when $P(A) > 0$, the conditional probability of B given A is

$$P(B|A) = P(A \text{ and } B) / P(A)$$

Statistics 528 - Lecture 17
Prof. Kate Calder 10