

## Midterm

**Problem 1.** In a half a page, discuss the advantages of a designed experiment over an observational study.

**Problem 2.** Recall from toxicological experiments that an important quantity of interest is the median lethal dose, or LD50. For the logistic model, the probability of death is related to dose through,

$$\log(\pi_i/(1 - \pi_i)) = \alpha + \beta \log d_i,$$

where  $\pi_i$  is the probability of death and  $d_i$  is the dose used in the  $i$ -th group;  $i = 1, \dots, k$ . Let  $(\hat{\alpha}, \hat{\beta})$  be the MLE of  $(\alpha, \beta)$  based on a binomial model in each group. (You do *not* have to know the formula for  $\hat{\alpha}$  and  $\hat{\beta}$ .)

(a) Derive a formula for LD50 in terms of  $\alpha$  and  $\beta$ .

(b) What is the MLE,  $\widehat{\text{LD50}}$ , of LD50?

**Problem 3.**

(a) How would you define the risk of a woman giving birth to a child with a birth defect, after exposure to contaminated drinking water during pregnancy?

(b) How would you define the relative risk of a birth defect, where “relative” here refers to exposure versus no exposure?

(c) Same question as in b), except replace “relative risk” with “odds ratio”.

**Problem 4.** Suppose a measurement  $X$  is made of a true concentration  $C$ . The “Method LOD” is defined as,

$$\text{LOD} = k \cdot \text{var}(X \mid C = 0),$$

Recall that the distribution of  $(X \mid C = 0)$  is lognormal  $(\mu, \sigma^2)$  if and only if

$$(Y \mid C = 0) \sim \text{normal}(\mu, \sigma^2),$$

where  $Y = \log X$ . Suppose we want to be sure that  $\Pr(X \geq \text{LOD} \mid C = 0) = 2.5\%$ . Assuming that  $X$  is lognormally distributed, find a formula for  $k$  in terms of  $\mu$  and  $\sigma$ , where you will need to know that the upper 2.5% point for a standard normal distribution is 1.96 and,

$$\text{var}(X \mid C = 0) = \exp(2\mu + \sigma^2)(\exp(\sigma^2) - 1).$$